

BALLOONING INSTABILITY IN POLAR CAPS OF ACCRETING NEUTRON STARS

C. LITWIN, EDWARD F. BROWN, AND R. ROSNER

Department of Astronomy & Astrophysics, The University of Chicago
Submitted to Astrophysical Journal

ABSTRACT

We assess the stability of Kruskal-Schwarzschild (magnetic Rayleigh-Taylor) type modes for accreted matter on the surface of a neutron star confined by a strong ($\gtrsim 10^{12}$ G) magnetic field. Employing the energy principle to analyze the stability of short-wavelength ballooning modes, we find that line-tying to the neutron star crust stabilizes these modes until the overpressure at the top of the neutron star crust exceeds the magnetic pressure by a factor $\sim 8(a/h)$, where a and h are respectively the lateral extent of the accretion region and the density scale height. The most unstable modes are localized within a density scale height above the crust. We calculate the amount of mass that can be accumulated at the polar cap before the onset of instability.

Subject headings: stars: neutron — pulsars — accretion, accretion disks — MHD — instabilities

1. INTRODUCTION

Despite recent progress on the spreading of accreted matter from a disk over an *unmagnetized* neutron star (Inogamov & Sunyaev 1999; Popham & Sunyaev 2000), there has been scarcely any effort directed at understanding how the magnetically funneled accretion stream of an accretion-powered X-ray pulsar spreads over the neutron star's surface. The presence of a strong magnetic field raises the possibility that the accreted matter is confined to the polar regions of the neutron star. This funneling and confinement of the accreted matter has been proposed (Joss & Li 1980) as the reason for the apparent stability of thermonuclear burning on strongly magnetized neutron stars. The presence of strong magnetic fields, with $B \gtrsim 10^{12}$ G, is deduced from cyclotron features in their spectrum (for reviews, see White et al. 1995; Heindl et al. 2000) and from regular torque reversals (Bildsten et al. 1997) that suggest the magnetospheric and co-rotation radii are similar. In neutron stars with weak magnetic fields (as inferred by the absence of pulsations or cyclotron features in the persistent emission), unstable burning is observed as type I X-ray bursts (Lewin et al. 1995). The funneling and confinement of the accreted matter by the magnetic field is believed to increase the local accretion rate to near- or super-Eddington magnitudes, for which the burning is stable to temperature perturbations (see Bildsten 1998, for a review).

Confinement of the accreted matter requires its overpressure be balanced by the magnetic field. Magnetostatic equilibria of this type were discussed by Hameury et al. (1983) and Brown & Bildsten (1998). In these papers, however, the stability of such equilibria was not addressed. This question is considered here.

If the magnetic field has an approximately potential dipole shape (as can be expected as long as the accreted plasma overpressure is low compared to the magnetic pressure), the force of gravity has a nonvanishing component perpendicular to the magnetic field. Such a configuration might be expected to be susceptible to Kruskal-Schwarzschild ("magnetic Rayleigh-Taylor") interchange instabilities (Kruskal & Schwarzschild 1954). Indeed, it

is easy to show that for characteristic parameters of polar cap plasmas the stabilizing effect of the magnetic curvature is small compared to the gravitational drive. Nevertheless, the line-tying in the neutron star crust eliminates interchange modes. Instead, ballooning modes can be unstable if the ratio of plasma overpressure to magnetic pressure $\Delta\beta$ is sufficiently large. This instability has been a subject of numerous investigations in the context of laboratory (e.g., Ohkawa & Kerst 1961; Coppi & Rosenbluth 1966; Connor et al. 1979; Freidberg 1987), solar (e.g., Hood 1986; De Bruyne & Hood 1989; Strauss & Longcope 1994), magnetospheric (e.g., McNutt et al. 1987; Hameiri et al. 1991) and astrophysical (e.g., Parker 1966; Shibata et al. 1989) plasmas. In the present context we find that this instability occurs when $\Delta\beta$ exceeds a/h where a is the polar cap radius and h is the density scale height near the neutron star crust. For such a large overpressure the equilibrium magnetic field is strongly distorted away from the potential field.

The structure of this paper is as follows. We begin by reviewing (§ 2) the conditions at the polar cap of an accreting neutron star. We then formulate, in section 3, the mathematical model which serves as the framework of our linear stability analysis. In section 4 we discuss the energy principle and derive a simple intuitive expression for the magnetohydrodynamic (MHD) potential energy, relevant for plasmas in the neutron star polar caps. We then use this expression to derive a criterion for stability of short-wavelength (in the direction transverse to the magnetic field) ballooning modes and discuss the stability of a plasma confined by an approximately potential, dipole magnetic field (§ 5). We then derive an approximate analytic expression for an equilibrium in which a large overpressure distorts the field away from the potential field (§ 6). Using this equilibrium we find the marginal stability criterion (§ 7), from which we compute the amount of mass that must be accreted in order to drive the instability (§ 8). We conclude with the discussion of our results in Section 9.

2. THE STRUCTURE OF POLAR CAPS

We are concerned with the stability of accreted matter on the polar cap of a strongly magnetized neutron star. Figure 1 provides a schematic of the situation. The collimated infalling matter approaches the neutron star at roughly free-fall velocity $v_{ff} \sim c$. At the near- or super-Eddington local accretion rates, some form of radiative shock is likely to form (Basko & Sunyaev 1976). The region we are interested in is far below where the infalling material decelerates and becomes incorporated into the neutron star atmosphere. The continual deposition of matter into the upper atmosphere pushes the underlying material deeper, and the downward flow velocity is $v \approx \dot{m}/\rho \ll c_s$, where $\dot{m}$ is the local accretion rate per unit area. The atmosphere is, to a high accuracy, in hydrostatic equilibrium. On time scales for matter to flow through the thickness of the atmosphere, the field lines co-move with the fluid (Brown & Bildsten 1998). Under these conditions we are justified in describing the structure of the accretion region with a magnetostatic approximation.

The polar cap is characterized by two length scales: its radius and the density scale height. Various estimates of the polar cap radius a (Lamb et al. 1973; Elsner & Lamb 1977; Arons & Lea 1980) suggest that $a \sim 10^5$ cm, and we shall adopt this value throughout the remainder of this paper. The density scale height h is much smaller than this value of a (see below). With these approximations, the mass and radius of the neutron star enter only through the surface gravity g , for which we use $g = 1.86 \times 10^{14}$ cm s $^{-2}$, which is the Newtonian value for a canonical neutron star of mass $M = 1.4M_\odot$ and a radius $R = 10$ km.




FIG. 1.— Overview of the accretion region.

Numerical solutions (Hameury et al. 1983; Brown & Bildsten 1998) of the magnetostatic equations show that the greatest distortion of the magnetic field occurs near the boundaries where the field lines are assumed to be tied. In the outermost layers of the neutron star, such a condition occurs at the neutron star crust where the ions form a Coulomb lattice. It is the rigidity of the crust that supports the superincumbent mass of the accreted matter in the polar cap and anchors the magnetic field lines. The electrostatic coupling is parameterized by

$$\Gamma = \frac{Z^2 e^2}{k_B T} \left(\frac{4\pi n_N}{3} \right)^{1/3}$$

$$\approx 1.1 \left(\frac{Z^2}{A^{1/3}} \right) \left(\frac{\rho}{10^8 \text{ g cm}^{-3}} \right)^{1/3} \frac{10^8 \text{ K}}{T} \quad (1)$$

where k_B is the Boltzmann constant, ρ is the density, and T , n_N , Ze , and A are, respectively, the temperature, the number density, the charge, and the atomic number of nuclei. For $\Gamma \gtrsim 1$, the ions form a liquid. For sufficiently large Γ , the ions form a crystalline lattice. The composition of the deep ocean and crust is unknown (see Schatz et al. 1999); for definiteness we take the ocean and crust to be composed of pure iron, and assume that solidification occurs for $\Gamma > 173$, in accordance with recent Monte-Carlo simulations (Farouki & Hamaguchi 1993). At this location, the crust of the neutron star can support a shear stress and therefore anchor the field lines. In the absence of a full calculation of the crust and magnetic field, we shall treat the crust [similarly as in previous works (Hameury et al. 1983; Brown & Bildsten 1998)] as a rigid, perfectly conducting surface to which the field lines are anchored (see, however, § 9).

As evident in equation (1), the location of the crust is determined by the temperature of the ocean. We assume that the burning of accreted hydrogen and helium is stable (this assumption can be checked *a posteriori* by finding the depth at which matter spreads). Stability of the H/He burning requires that the accretion rate be locally super-Eddington (see Bildsten 1998); for the remainder of this work we shall use the local Eddington rate as our fiducial value. In this case, the temperature in the ocean is set by the nuclear flux released from burning of the accreted fuel to iron peak elements (about 8 MeV per nucleon) and is $\gtrsim 5 \times 10^8$ K (Brown & Bildsten 1998; Schatz et al. 1999).

For the density and temperature of interest, the electrons are degenerate, i.e., $k_B T < E_F - m_e c^2$, where $E_F = m_e c^2 [1 + (3\pi^2 n_e)^{2/3} (\hbar/m_e c)^2]^{1/2}$ is the electron Fermi energy, and m_e and n_e are the electron mass and density, respectively. The principal contribution to pressure p is the electron pressure p_e and therefore to a good approximation $p \approx p_e \approx n_e E_F / 4 \propto \rho^\gamma$ where $\gamma = 4/3$ and the proportionality constant is only a function of the ion charge-to-mass ratio Z/A . For a plasma of uniform composition this constant is independent of both time and space. We shall use this approximation in the next section.

Using this approximate equation of state, we write the density scale height $h = \gamma p / \rho g$ at the top of the crust in terms of Γ ,

$$\begin{aligned} h &\approx \gamma \left(\frac{9\pi}{256} \right)^{1/3} \frac{1}{Z^{2/3} A} \frac{N_A k_B T}{g} \frac{\Gamma}{\alpha_F} \\ &= 6.90 \times 10^3 \text{ cm} \\ &\times \left(\frac{26}{Z} \right)^{2/3} \left(\frac{56}{A} \right) \left(\frac{T}{5 \times 10^8 \text{ K}} \right) \left(\frac{\Gamma}{173} \right), \quad (2) \end{aligned}$$

where $\gamma = 4/3$, $N_A = 6.02 \times 10^{23}$ g $^{-1}$ is Avogadro's constant and $\alpha_F = e^2 / \hbar c$.

3. MODEL

We model the polar cap as the region of the neutron star atmosphere and ocean that is bounded by the same magnetic flux surface within which accretion occurs. This region is characterized by a somewhat higher pressure than in the surrounding matter, due to the weight of

the accreted material. The plasma in the polar cap and in the surrounding region will be treated as a perfectly conducting fluid of uniform composition, bounded by a perfectly conducting, rigid surface (the top of the crust) and immersed in a magnetic field. The magnetic field lines are assumed tied to the crust; the intersection point of the field line with the top of the crust will be referred to as the footpoint. Since the density scale height of the plasma in the polar cap is much smaller than the stellar radius we shall consider the field lines as semi-infinite and having only one footpoint. In the equilibrium, the plasma and the immersing magnetic field are assumed to be axisymmetric with the axis of symmetry coinciding with the magnetic axis; the magnetic field is assumed to be untwisted.

We describe the plasma by the single-fluid MHD theory. In the presence of the gravitational potential U and magnetic field $\mathbf{B}$ the momentum balance equation for a plasma of density ρ and pressure p , moving with velocity $\mathbf{v}$, is

$$\rho \frac{d\mathbf{v}}{dt} = \frac{1}{c} \mathbf{J} \times \mathbf{B} - \nabla p - \rho \nabla U. \quad (3)$$

While Ohm's law is modified by gravity, the induction equation is unaffected (cf. Bernstein et al. (1958)), so that

$$\frac{\partial \mathbf{B}}{\partial t} = \nabla \times \mathbf{v} \times \mathbf{B}. \quad (4)$$

The above equations are supplemented by Ampère's law,

$$\mathbf{J} = \frac{c}{4\pi} \nabla \times \mathbf{B}, \quad (5)$$

the continuity equation,

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{v}) = 0, \quad (6)$$

and an equation of state which we assume to be adiabatic:

$$\frac{d}{dt} p \rho^{-\gamma} = 0. \quad (7)$$

As mentioned in §1, the primary contribution to the plasma pressure in the region of interest is due to degenerate, relativistic electrons ($\gamma = 4/3$). Therefore, in region of uniform composition, $p \rho^{-\gamma}$ is, to a high accuracy, constant in space. For the assumptions in § 2, $\epsilon \equiv d \ln p \rho^{-1/\gamma} / d \ln p \approx 3/8 p_{\text{ion}} / p_e = (3/2Z)(k_B T / E_F) \approx 4 \times 10^{-4}$. Thus

$$p \rho^{-\gamma} = \text{const.}, \quad (8)$$

to a very good approximation.

The force balance in a static equilibrium is given by

$$\frac{1}{c} \mathbf{J} \times \mathbf{B} = \nabla p + \rho \nabla U, \quad (9)$$

from which it follows that

$$\nabla p + \rho \nabla U = \rho \nabla F, \quad (10)$$

where the function F is a function constant on the field lines

$$\mathbf{B} \cdot \nabla F = 0. \quad (11)$$

By setting the gravitational potential to zero at the top of the crust, F is determined by the sound speed $c_s = \sqrt{\gamma p / \rho}$ there, $F = c_{s0}^2 / (\gamma - 1)$.

An untwisted, axisymmetric equilibrium magnetic field can be represented in the form

$$\mathbf{B} = \nabla \psi \times \nabla \theta \quad (12)$$

where ψ is the flux function and θ is the toroidal angle. It is straightforward to show with the aid of Ampère law that in such a geometry the current is perpendicular to the magnetic field: $\mathbf{J} \cdot \mathbf{B} = 0$. Eq. (11) implies that F is a function of the flux function only: $F = F(\psi)$.

Upon introducing cylindrical coordinates (r, θ, z) with the z -direction along the axis of symmetry, the normal (i.e., parallel to $\nabla \psi$) component of Eq. (10) yields the Grad-Shafranov-like equation

$$r \frac{\partial}{\partial r} \frac{1}{r} \frac{\partial \psi}{\partial r} + \frac{\partial^2 \psi}{\partial z^2} = -4\pi r^2 \rho(\psi, z) \frac{dF}{d\psi}. \quad (13)$$

with ρ being determined by equation (10).

In the limit $a \ll R$ the curvature of the stellar surface can be neglected on the scale of the accretion region. In this approximation $U = gz$, so that

$$\rho(\psi, z) = \rho_0(\psi) \left[1 - \alpha \frac{z}{h(\psi)} \right]^{1/\alpha} \quad (14)$$

where $z = 0$ corresponds to the top of the crust, $\alpha \equiv \gamma - 1$, ρ_0 is the plasma density at $z = 0$, $h \equiv -(d \ln \rho / dz)^{-1}|_{z=0} = \alpha F / g$ is the density scale height at the footpoints and g is the gravitational acceleration. Note that ρ_0 and h are functions of ψ .

4. ENERGY PRINCIPLE

We now address the question of the linear stability of the polar cap plasma in the approximation described in the previous section. We base our analysis on the MHD energy principle (Bernstein et al. 1958). In the present section we discuss the energy principle as applicable to the polar cap plasmas, generalizing the “intuitive” form of the energy principle (Furth et al. 1965; Greene & Johnson 1968) to include the effects of a strong gravitational field. In our discussion we follow the approach of Freidberg (1987).

We start by considering the linearized momentum balance equation for a static equilibrium

$$\rho \frac{\partial^2 \boldsymbol{\xi}}{\partial t^2} = \mathbf{F}(\boldsymbol{\xi}) \equiv \frac{1}{c} \tilde{\mathbf{J}} \times \mathbf{B} + \frac{1}{c} \mathbf{J} \times \tilde{\mathbf{B}} - \nabla \tilde{p} - \tilde{\rho} \nabla U \quad (15)$$

where $\boldsymbol{\xi}$ is the plasma displacement and the gravitational potential is unperturbed. Here and in the following the tilde denotes perturbed quantities while symbols without it stand for the equilibrium quantities. Perturbed current $\tilde{\mathbf{J}}$, magnetic field $\tilde{\mathbf{B}}$, density $\tilde{\rho}$, and pressure $\tilde{p}$ are respectively given by Ampère's law, the induction equation, the continuity equation, and the equation of state:

$$\tilde{\mathbf{J}} = \frac{c}{4\pi} \nabla \times \tilde{\mathbf{B}} \quad (16)$$

$$\tilde{\mathbf{B}} = \nabla \times (\boldsymbol{\xi} \times \mathbf{B}) \quad (17)$$

$$\tilde{\rho} = -\nabla \cdot (\rho \boldsymbol{\xi}) \quad (18)$$

$$\tilde{p} = -\gamma p \nabla \cdot \boldsymbol{\xi} - \boldsymbol{\xi} \cdot \nabla p. \quad (19)$$

We shall assume that the plasma boundary (the crust) is rigid and perfectly conducting, so that $\boldsymbol{\xi}$ and $\tilde{\mathbf{B}}$ vanish there.

The second-order potential energy is given by (Bernstein et al. 1958)

$$\begin{aligned} \delta W &= -\frac{1}{2} \int d\tau \boldsymbol{\xi} \cdot \mathbf{F}(\boldsymbol{\xi}) \\ &= -\frac{1}{2} \int d\tau \boldsymbol{\xi} \cdot \left\{ \frac{1}{c} \tilde{\mathbf{J}} \times \mathbf{B} + \frac{1}{c} \mathbf{J} \times \tilde{\mathbf{B}} \right. \\ &\quad \left. + \nabla(\gamma p \nabla \cdot \boldsymbol{\xi} + \boldsymbol{\xi} \cdot \nabla p) + \nabla \cdot (\rho \boldsymbol{\xi}) \nabla U \right\} \end{aligned} \quad (20)$$

where $d\tau$ denotes integration over the plasma volume. We now transform the above expression for the potential energy into a form that allows a physical interpretation. This is analogous to what Furth et al. (1965) and Greene & Johnson (1968) have done in the absence of gravitational field. In order to derive it, we follow the standard approach [e.g., Freidberg (1987)], with appropriate modifications for the presence of a strong gravitational field.

Integrating by parts and dropping surface terms (which vanish because of the assumed boundary conditions), and exploiting the equilibrium relation given by Eq. (10), one obtains, after some straightforward algebra,

$$\begin{aligned} \delta W &= \frac{1}{2} \int d\tau \left\{ \gamma p (\nabla \cdot \boldsymbol{\xi} - 2\boldsymbol{\kappa}_g \cdot \boldsymbol{\xi})^2 + \epsilon p (\boldsymbol{\xi} \cdot \nabla \ln p)^2 + \frac{\tilde{\mathbf{B}}^2}{4\pi} \right. \\ &\quad \left. - \boldsymbol{\xi} \cdot \left[\frac{1}{c} \mathbf{J} \times \tilde{\mathbf{B}} + 2\boldsymbol{\kappa}_g (\rho \boldsymbol{\xi} \cdot \nabla F) + \nabla (\rho \boldsymbol{\xi} \cdot \nabla F) \right] \right\}. \end{aligned} \quad (21)$$

where $\boldsymbol{\kappa}_g \equiv (\rho/\gamma p) \nabla U/2$. The second term on the RHS, proportional to the Schwarzschild discriminant, can be neglected for highly degenerate plasmas in strong magnetic fields because of the smallness of parameter $\epsilon = d \ln \rho p^{-1/\gamma} / d \ln p$. We shall omit this term in the subsequent analysis (see, however, discussion of finite ϵ effects in §9).

The last two terms in the square bracket on the RHS of Eq. (21) can be expanded and recombined to yield an alternative form of the energy integral

$$\begin{aligned} \delta W &= \frac{1}{2} \int d\tau \left\{ \frac{\tilde{\mathbf{B}}^2}{4\pi} + \gamma p (\nabla \cdot \boldsymbol{\xi} - 2\boldsymbol{\kappa}_g \cdot \boldsymbol{\xi})^2 \right. \\ &\quad \left. - \boldsymbol{\xi} \cdot \left[\frac{1}{c} \mathbf{J} \times \tilde{\mathbf{B}} + \frac{\rho^2}{\gamma p} (\boldsymbol{\xi} \cdot \nabla F) \rho \nabla F + \rho \nabla (\boldsymbol{\xi} \cdot \nabla F) \right] \right\}. \end{aligned} \quad (22)$$

Since $\mathbf{B} \cdot \nabla F = 0$ and

$$\frac{1}{c} \mathbf{B} \cdot \mathbf{J} \times \tilde{\mathbf{B}} = -\rho \mathbf{B} \cdot \nabla (\boldsymbol{\xi} \cdot \nabla F), \quad (23)$$

it is clear that the parallel (to $\mathbf{B}$) component of the expression in the square bracket in Eq. (22), and hence also in Eq. (21), vanishes. Thus the energy integral can be expressed in the form

$$\begin{aligned} \delta W &= \frac{1}{2} \int d\tau \left\{ \frac{\tilde{\mathbf{B}}^2}{4\pi} + \gamma p (\nabla \cdot \boldsymbol{\xi} - 2\boldsymbol{\kappa}_g \cdot \boldsymbol{\xi})^2 \right. \\ &\quad \left. - \boldsymbol{\xi}_\perp \cdot \left[\frac{1}{c} \mathbf{J} \times \tilde{\mathbf{B}} + 2(\rho \boldsymbol{\xi}_\perp \cdot \nabla F) \boldsymbol{\kappa}_g + \nabla (\rho \boldsymbol{\xi}_\perp \cdot \nabla F) \right] \right\} \end{aligned} \quad (24)$$

Next, splitting up $\tilde{\mathbf{B}}$ into transverse and parallel components, $\tilde{\mathbf{B}} = \tilde{\mathbf{B}}_\perp + \tilde{B}_\parallel \hat{\mathbf{b}}$ ($\hat{\mathbf{b}} \equiv \mathbf{B}/B$), we find

$$\boldsymbol{\xi}_\perp \cdot \mathbf{J} \times \tilde{\mathbf{B}} = J_\parallel \boldsymbol{\xi}_\perp \cdot \hat{\mathbf{b}} \times \tilde{\mathbf{B}} + \tilde{B}_\parallel \boldsymbol{\xi}_\perp \cdot \mathbf{J} \times \hat{\mathbf{b}} \quad (25)$$

$$\tilde{B}_\parallel = -B(\nabla \cdot \boldsymbol{\xi}_\perp + 2\boldsymbol{\kappa}_c \cdot \boldsymbol{\xi}_\perp) + \frac{4\pi\rho}{B} \boldsymbol{\xi}_\perp \cdot \nabla F, \quad (26)$$

where $\boldsymbol{\kappa}_c \equiv \hat{\mathbf{b}} \cdot \nabla \hat{\mathbf{b}}$ is the field line curvature. With these expressions the potential energy can be expressed in the following, “intuitive” form

$$\begin{aligned} \delta W &= \frac{1}{2} \int d\tau \left\{ \frac{\tilde{\mathbf{B}}_\perp^2}{4\pi} + \frac{B^2}{4\pi} (\nabla \cdot \boldsymbol{\xi}_\perp + 2\boldsymbol{\kappa}_c \cdot \boldsymbol{\xi}_\perp)^2 \right. \\ &\quad \left. + \gamma p (\nabla \cdot \boldsymbol{\xi} - 2\boldsymbol{\kappa}_g \cdot \boldsymbol{\xi})^2 - \frac{1}{c} J_\parallel \boldsymbol{\xi}_\perp \times \hat{\mathbf{b}} \cdot \tilde{\mathbf{B}} \right. \\ &\quad \left. - 2\boldsymbol{\kappa}_c \cdot \boldsymbol{\xi}_\perp \rho (\boldsymbol{\xi}_\perp \cdot \nabla F) \right\}. \end{aligned} \quad (27)$$

where $\boldsymbol{\kappa} \equiv \boldsymbol{\kappa}_c + \boldsymbol{\kappa}_g$. The first term is the energy corresponding to field line bending, the second is the field compression energy, the third is the energy of plasma compression and buoyancy, the fourth one is responsible for the kink instability, and the last one drives the curvature and gravitational instabilities. The above expression is a generalization of the result of Furth et al. (1965) and Greene & Johnson (1968) to the case of a degenerate plasma in a strong gravitational field. The above expression reduces to these earlier results if the density scale height is much longer than the plasma length, i.e., if $U \ll c_s^2$.

Thus far we have not taken into account the geometry of polar caps and the derived energy principle (eq. [27]) applies to an arbitrary configuration of plasmas satisfying the equation of state (7). We shall now perform a reduction of the energy principle as applicable to the configuration of our interest.

First we note that the parallel displacement $\xi_\parallel$ appears only in the nonnegative compressive term. Therefore this term can be minimized, in a standard manner, by considering the perturbation of the compressive energy due to a perturbation of $\xi_\parallel$. The minimum is found for the displacement satisfying the Euler-Lagrange equation which can be expressed in the form

$$\mathbf{B} \cdot \nabla \left[p^{1-\frac{1}{\gamma}} (\nabla \cdot \boldsymbol{\xi} - 2\boldsymbol{\kappa}_g \cdot \boldsymbol{\xi}) \right] = 0 \quad (28)$$

Integrating along the field line and imposing the line-tying boundary condition ($\xi_\parallel[0] = 0$) yields

$$\xi_\parallel = \frac{B}{p^{\frac{1}{\gamma}}} \int_0^l \frac{dl'}{B} p^{\frac{1}{\gamma}} \left[p^{\frac{1}{\gamma}-1} A + \nabla \cdot \boldsymbol{\xi}_\perp - 2\boldsymbol{\kappa}_g \cdot \boldsymbol{\xi}_\perp \right] \quad (29)$$

where l is the distance along the field line. The integration constant A must be determined from the integrability condition. Note that $p, \rho \rightarrow 0$ as $z \rightarrow z_0 \equiv h/(\gamma-1)$ (cf. equations [14] and [8]). Therefore, for the kinetic energy of parallel motion to be finite, (i.e., $\int d\tau \rho \xi_\parallel^2 < \infty$), this requires that

$$A = - \frac{\int_0^{l_0} \frac{dl'}{B} p^{\frac{1}{\gamma}} (\nabla \cdot \boldsymbol{\xi}_\perp - 2\boldsymbol{\kappa}_g \cdot \boldsymbol{\xi}_\perp)}{\int_0^{l_0} \frac{dl'}{B} p^{\frac{2}{\gamma}-1}} \quad (30)$$

where $l_0 \equiv l(z_0)$. This implies, in particular, that $\xi_\parallel(z_0) = 0$. Since

$$\nabla \cdot \boldsymbol{\xi} - 2\boldsymbol{\kappa}_g \cdot \boldsymbol{\xi} = p^{\frac{1}{\gamma}-1}A \quad (31)$$

and $A \neq 0$, the compressional energy cannot be minimized to zero. This is analogous to that which occurs in flux tubes line-tied at both ends (cf. Hameiri et al 1991) and is, in fact, caused by the effective line-tying in the low density region.

Setting $J_{\parallel} = 0$, as appropriate for an untwisted axisymmetric field, the potential energy becomes

$$\begin{aligned} \delta W = \frac{1}{2} \int d\tau \left\{ \frac{\tilde{\mathbf{B}}_{\perp}^2}{4\pi} + \frac{\mathbf{B}^2}{4\pi} (\nabla \cdot \boldsymbol{\xi}_{\perp} + 2\boldsymbol{\kappa}_c \cdot \boldsymbol{\xi}_{\perp})^2 \right. \\ \left. + \gamma p^{\frac{2}{\gamma}-1} \left[\frac{\int_0^{l_0} \frac{dl'}{B} p^{\frac{1}{\gamma}} (\nabla \cdot \boldsymbol{\xi}_{\perp} - 2\boldsymbol{\kappa}_g \cdot \boldsymbol{\xi}_{\perp})}{\int_0^{l_0} \frac{dl'}{B} p^{\frac{2}{\gamma}-1}} \right]^2 \right. \\ \left. - 2\rho(\boldsymbol{\xi}_{\perp} \cdot \nabla F) \boldsymbol{\kappa} \cdot \boldsymbol{\xi}_{\perp} \right\}. \quad (32) \end{aligned}$$

5. BALLOONING INSTABILITY

We now limit our attention to the short wavelength (in the direction perpendicular to the magnetic field) modes. Following the method of Freidberg (1987), we perform the reduction of the intuitive form of the energy principle (eq. [32]).

We represent the displacement in the eikonal approximation

$$\boldsymbol{\xi}_{\perp} = \boldsymbol{\eta}_{\perp} e^{iS} + c.c. \quad (33)$$

where *c.c.* stands for complex conjugate; $\boldsymbol{\eta}_{\perp}$ is assumed to be slowly varying while the eikonal S is assumed to vary rapidly, i.e.,

$$\mathbf{k}_{\perp} = \nabla S \quad (34)$$

is so large that $k_{\perp} a \gg 1$, where a is the equilibrium length scale.

Note that $\mathbf{B} \cdot \nabla S = 0$. Consequently

$$\tilde{\mathbf{B}}_{\perp} = e^{iS} [\nabla \times (\boldsymbol{\eta}_{\perp} \times \mathbf{B})]_{\perp} + c.c. \quad (35)$$

In the limit of $k_{\perp} \rightarrow \infty$, the dominant terms in δW (Eq.[32]) are those proportional to $(\nabla \cdot \boldsymbol{\xi}_{\perp})^2 \approx |\mathbf{k}_{\perp} \cdot \boldsymbol{\eta}_{\perp}|^2$ (second and third term on the RHS of eq. [32]). Therefore the minimization of the potential energy, to lowest order in $1/k_{\perp} a$, requires that $\mathbf{k}_{\perp} \cdot \boldsymbol{\eta}_{\perp} = 0$, which implies that for the zeroth-order displacement

$$\boldsymbol{\eta}_{\perp} \approx \frac{X}{B} \hat{\mathbf{b}} \times \mathbf{k}_{\perp}. \quad (36)$$

where X is a scalar function. To the next order, the first-order displacement can be chosen so that the sum of two positive terms involving $\nabla \cdot \boldsymbol{\xi}_{\perp}$ (second and third term on the RHS of eq. [32]) is minimized. Since we are interested in the limit of $\beta \equiv 8\pi p/B^2 \gg 1$, we can instead minimize only the second of these terms. A simple choice of a minimizing displacement is

$$\nabla \cdot \boldsymbol{\xi}_{\perp} = 2\boldsymbol{\kappa}_g \cdot \boldsymbol{\xi}_{\perp}. \quad (37)$$

For this choice, $\xi_{\parallel} = 0$ as well as the third term on the RHS of eq. (32) vanish so that

$$\begin{aligned} \delta W = \frac{1}{2} \int d\tau \left\{ \frac{\tilde{\mathbf{B}}_{\perp}^2}{4\pi} + \frac{\mathbf{B}^2}{\pi} (\boldsymbol{\kappa} \cdot \boldsymbol{\xi}_{\perp})^2 \right. \\ \left. - 2\rho(\boldsymbol{\xi}_{\perp} \cdot \nabla F)(\boldsymbol{\kappa} \cdot \boldsymbol{\xi}_{\perp}) \right\}. \quad (38) \end{aligned}$$

From equation (36) it follows that

$$\tilde{\mathbf{B}}_{\perp} = e^{iS} (\nabla X \times \hat{\mathbf{b}})_{\perp} + c.c. = (\hat{\mathbf{b}} \cdot \nabla X) \hat{\mathbf{b}} \times \mathbf{k}_{\perp} + c.c. \quad (39)$$

Consequently, one finds from equation (38), exploiting the hermiticity of the operator $\mathbf{F}$ (Bernstein et al. 1958), that

$$\begin{aligned} \delta W = \frac{1}{4\pi} \int d\tau \left\{ k_{\perp}^2 |\hat{\mathbf{b}} \cdot \nabla X|^2 + 4(\boldsymbol{\kappa} \times \hat{\mathbf{b}} \cdot \mathbf{k}_{\perp})^2 |X|^2 \right. \\ \left. - \frac{8\pi\rho}{B^2} (\boldsymbol{\kappa} \times \hat{\mathbf{b}} \cdot \mathbf{k}_{\perp})(\nabla F \times \hat{\mathbf{b}} \cdot \mathbf{k}_{\perp}) |X|^2 \right\}. \quad (40) \end{aligned}$$

For an axisymmetric system, $S = m\theta + \hat{S}(\psi, l)$ where l is the distance along the field line. In the absence of a toroidal field B_{θ} , $\mathbf{B} \cdot \nabla S = 0$ implies that $\hat{S} = \hat{S}(\psi)$. Thus

$$\mathbf{k}_{\perp} = k_{\theta} \hat{\boldsymbol{\theta}} + k_n \hat{\mathbf{n}} \quad (41)$$

where $k_{\theta} = m/r$ and $\hat{\mathbf{n}} = \nabla\psi/|\nabla\psi| = \nabla\psi/rB$ is the unit vector normal to the flux surface. Noting that

$$\hat{\mathbf{b}} \times \mathbf{k}_{\perp} = -k_{\theta} \hat{\mathbf{n}} + k_n \hat{\boldsymbol{\theta}} \quad (42)$$

$$\hat{\mathbf{b}} \times \mathbf{k}_{\perp} \cdot \nabla F = -k_{\theta} r B \frac{dF}{d\psi}, \quad (43)$$

the potential energy can be put in form

$$\begin{aligned} \delta W = \frac{1}{4\pi} \int d\tau \left\{ (k_{\theta}^2 + k_n^2) \left| \frac{\partial X}{\partial l} \right|^2 \right. \\ \left. + 4k_{\theta}^2 \kappa_n (\kappa_n - \frac{2\pi r \rho F'}{B}) |X|^2 \right\} \quad (44) \end{aligned}$$

($F' = dF/d\psi$). Minimization with respect to k_n is achieved by setting $k_n = 0$.

The integrand in Eq. (44) involves derivatives of X only along the magnetic field but not derivatives of ψ . Hence ψ can be treated as a parameter and the potential energy can be expressed in the form

$$\delta W = \int d\psi \delta W(\psi), \quad (45)$$

where

$$\begin{aligned} \delta W(\psi) = \frac{m^2}{4\pi} \int \frac{dl}{r^2 B} \left\{ \left| \frac{\partial X}{\partial l} \right|^2 \right. \\ \left. + 4\kappa_n (\kappa_n - \frac{2\pi r \rho F'}{B}) |X|^2 \right\}. \quad (46) \end{aligned}$$

Equation (45) implies that $\delta W(\psi) < 0$ for some ψ is both a *necessary* and *sufficient* condition for instability (Bernstein et al. 1958).

Let us first address the question of stability of a plasma confined by a stellar dipole magnetic field. We approximate F' by $\Delta F/\psi_a$ where $\psi_a = a^2 B_0/2$ and $\Delta F = \Delta p_0/\rho_0$, with Δp_0 and ρ_0 being, respectively, the overpressure and density at the top of the crust. Since $\kappa_{gn} = (\partial U/\partial n)/2c_{s0}^2 \sim -\hat{b}_r/2h$, $\kappa_{cn} \sim \hat{b}_r/R$, $\hat{b}_r \sim r/R$ (for a dipole field) and the density scale height $h = c_{s0}^2/g$ is much smaller than the neutron star radius R , the second term on the RHS of Eq. (46) is destabilizing, if $\Delta\beta \gtrsim a^2/hR$.

For the interchange modes $\partial X/\partial l = 0$. Because of line tying, however, $X(0) = 0$ with $l = 0$ being chosen to correspond to the top of the crust. Expanding into Taylor series, we can approximate $X \sim l$ for $l \lesssim h$. From the structure of the integral in Eq. (46) we expect that the most unstable displacement vanishes for $l \gg h$.

We shall now estimate the integral in Eq. (46). To lowest order in a/R , where a is the radius of the polar cap, $U = c_s^2 l/h$. We then find that $\delta W(\psi_a)$ becomes negative only if

$$\Delta\beta \gtrsim \frac{R}{h} \quad (47)$$

where $\Delta\beta = 8\pi\Delta p_0/B_0^2$. The above equation implies that for a low overpressure ($\Delta\beta \lesssim 1$), the plasma confined by the dipole magnetic field of a neutron star is stable with respect to ballooning modes. Equation (47) indicates that the plasma may become unstable at high $\Delta\beta \gg 1$. For such a high plasma overpressure, however, the equilibrium magnetic field is distorted away from the dipole field and has to be computed self-consistently. We consider this case in the next section.

6. A SIMPLE HIGH- $\Delta\beta$ EQUILIBRIUM

In the limit $a \ll R$ the dipole magnetic field in the polar cap is well approximated by a uniform field. Let us therefore assume that prior to the loading of the accretion column with plasma, the magnetic field $\mathbf{B} = B_0 \hat{z}$. We shall assume that at $z = 0$ the flux at the is unaffected by the accretion, i.e., $\psi(r, 0) = r^2 B_0/2$. The loading is assumed to occur on flux surfaces with $\psi \leq \psi_a \equiv a^2 B_0/2$.

Until now, the overpressure profile at the base of the accretion region, which determines the function F , has not been specified. As a simple example we shall consider the following form of F ,

$$F(\psi) = F_{ext} + \Delta F \left[1 - \left(\frac{\psi}{\psi_a} \right)^2 \right]^{1+\varepsilon} \quad (48)$$

for $\psi \leq \psi_a \equiv a^2 B_0/2$, with $\varepsilon \ll 1$; F is a constant F_{ext} for $\psi > \psi_a$. The gradient of F is approximately linear for $\psi < \psi_a(1 - \varepsilon)$ and goes smoothly to zero at $\psi = \psi_a$. In the following we shall restrict our attention to the case of an overpressure Δp_0 that is small compared to the total pressure p_0 , so that $\Delta F = \Delta p_0/\rho_0$. In the same approximation, the scale height h is, to lowest order, independent of ψ .

For such a pressure profile, the Grad-Shafranov equation (Eq. [11]) inside the accretion column [$\psi < \psi_a(1 - \varepsilon)$] becomes approximately linear,

$$r \frac{\partial}{\partial r} \left(\frac{1}{r} \frac{\partial \psi}{\partial r} \right) + \frac{\partial^2 \psi}{\partial z^2} = \frac{4\Delta\beta}{a^2} \left(\frac{r}{a} \right)^2 \left(1 - \alpha \frac{z}{h} \right)^{\frac{1}{\alpha}} \psi. \quad (49)$$

We anticipate that because $\Delta\beta \gg 1$ and $h \ll a$, the radial excursion of the flux surfaces $\Delta r = \int dl \hat{b}_r$ ($\lesssim h$) will be small compared to r . We can therefore neglect the r -dependence on the RHS of Eq. (49) and treat $\chi \equiv r/a$ as a parameter. With these approximations, we can then solve Eq. (49) by separation of variables,

$$\psi(r, z) = \psi_0 \chi^2 f(\zeta)|_{\chi=r/a, \zeta=z/h}, \quad (50)$$

where f satisfies the equation

$$\frac{d^2 f}{d\zeta^2} = \lambda^2 \chi^2 (1 - \alpha\zeta)^{\frac{1}{\alpha}} f \quad (51)$$

($\lambda \equiv 2h\sqrt{\Delta\beta}/a$). We solve this equation subject to the boundary conditions $f(0) = 1$ and $f'(\alpha^{-1}) = 0$. The first of these conditions expresses normalization; the second follows from the requirement that the field be vertical in the region of vanishing overpressure. By changing variables, we can transform the above equation into a modified Bessel equation which has the solution

$$f(\zeta) = u^\nu(\zeta) \{ \mathcal{A} I_{-\nu}[u(\zeta)] + \mathcal{B} I_\nu[u(\zeta)] \}, \quad (52)$$

where $u(\zeta) = 2\lambda\chi(1 - \alpha\zeta)^{\frac{1}{2\alpha}}/(1 + 2\alpha)$, $\nu = \alpha/(1 + 2\alpha) = (\gamma - 1)/(2\gamma - 1)$, $I_{\pm\nu}$ are the modified Bessel functions, and $\mathcal{A}$ and $\mathcal{B}$ are the integration constants. From the condition $f'(\alpha^{-1}) = 0$ it follows that $\mathcal{B} = 0$; the boundary condition $f(0) = 1$ then implies that

$$f(\zeta) = (1 - \alpha\zeta)^{\frac{1}{2}} \frac{I_{-\nu} \left[\frac{2\lambda\chi}{1+2\alpha} (1 - \alpha\zeta)^{\frac{1}{2\alpha}} \right]}{I_{-\nu} \left(\frac{2\lambda\chi}{1+2\alpha} \right)}. \quad (53)$$

An example of flux surfaces given by equations (50) and (53) is shown in Figure 2. We see that even for relatively large λ , the radial excursion Δr of flux surfaces is small compared to r , in accord with our assumptions.




FIG. 2.— Poloidal cross-sections of flux surfaces for $\lambda = 1$ (solid) and $\lambda = 0.5$ (dashed), with $a/h = 10$ (r and z are in units of a).

Let us now consider the case of $a/h \lesssim \Delta\beta \ll a^2/h^2$ which, as we shall see, is valid at the onset of instability. In this case, $\lambda \ll 1$ and we can expand the Bessel functions into a power series. In this approximation we find that

$$\psi(r, z) = \psi_0 \frac{r^2}{a^2} \left[1 - \lambda^2 \frac{r^2}{a^2} \frac{1 - (1 - \alpha z/h)^{2+\frac{1}{\alpha}}}{(1 + \alpha)(1 + 2\alpha)} \right]. \quad (54)$$

With the aid of the above expression we can now demonstrate the self-consistency of our earlier assumptions. We find that the first term on the LHS of Eq. (49) is smaller than the second one as long as $\rho/\rho_0 > (2h/a)^2$. Thus for $h \ll a$ our solution is indeed self-consistent in the region of plasma where the most unstable displacement is expected to be localized (see §7).

From (54) the magnetic field components are

$$B_r \approx \mu \left(\frac{\rho}{\rho_0} \right)^\gamma B_0 \quad (55)$$

$$B_z \approx B_0 \quad (56)$$

where $\mu = 2\Delta\beta h x^3/\gamma a$. From the above it follows that the angle ϕ between the magnetic and gravitational fields is given by

$$\tan \phi \approx \mu \left(1 - \alpha \frac{z}{h} \right)^{\gamma/\alpha}. \quad (57)$$

From equation (57) one sees explicitly that the field line distortion becomes large ($\tan \phi \gtrsim 1$) when $\mu \gtrsim 1$, i.e., $\Delta\beta \gtrsim a/h$, as originally observed by Hameury et al. (1983). This is, in fact, a simple consequence of balancing overpressure $\Delta p/a$ with magnetic tension $\kappa B^2/4\pi$ since $\kappa h \sim 1$ when the field distortion becomes large.

7. CRITERION FOR THE ONSET OF INSTABILITY

We now use the equilibrium found in the previous section to evaluate the energy integral. Let us first change the integration variable in Eq. (46) from l to z , such that $dl = dz/\hat{b}_z$. For the nearly parabolic pressure profile considered earlier,

$$\delta W(\psi) = \frac{m^2}{8\pi} \int \frac{dz}{r^2 B_z} \left\{ \hat{b}_z^2 \left| \frac{\partial X}{\partial z} \right|^2 + 4\kappa_n \left(\kappa_n + \frac{\gamma\mu B_0}{2h} \frac{\rho}{\rho_0} \right) |X|^2 \right\}. \quad (58)$$

In the approximation of Eq. (54), we find that to lowest order in λ

$$\kappa_n \approx -(1 + 2\gamma \hat{b}_z^2) \left(\frac{\rho_0}{\rho} \right)^\alpha \frac{\hat{b}_r}{2h}. \quad (59)$$

Changing the integration variable to $s = \tan \phi$, as given by equation (57), we then find, with the aid of (55), (56), (57) and (59),

$$\delta W(\psi) = \frac{\mu^{1-1/\gamma} m^2}{8\pi r^2 \gamma h B_0} \int_0^\mu ds \frac{s^{1/\gamma}}{1+s^2} \left\{ \gamma^2 \left| \frac{\partial X}{\partial s} \right|^2 - (\gamma-1) \frac{(s^2 - \mu_0^2)(s^2 + 2\gamma + 1)}{(s^2 + 1)^2} |X|^2 \right\}, \quad (60)$$

where $\mu_0 = \sqrt{(\gamma+1)/(\gamma-1)}$. We observe that the integrand in equation (60) is positive for $s \geq \mu_0$. Consequently, the most unstable displacement is expected to vanish for $s \geq \mu_0$ which implies the instability is localized to the region where

$$\rho \geq \left(\frac{\mu_0}{\mu} \right)^{\frac{1}{\gamma}} \rho_0. \quad (61)$$

Since the line-tying in the crust requires that $X(s = \mu) = 0$, we therefore choose as a test function

$$X(s) = X_0 \sin \left(\pi \frac{s - \mu_0}{\mu - \mu_0} \right)$$

For this test function and $\gamma = 4/3$, as appropriate for a plasma in the vicinity of the crust, we find that $\delta W(\psi) < 0$ for $\mu > 11.67$ for which it follows that

$$\Delta\beta > 7.8 \frac{a}{h}. \quad (62)$$

is a sufficient condition for instability. For $\mu = 11.67$, equation (61) implies that the instability is localized to within 0.93 scale height from the crust.

The instability criterion (62) is obtained for the equilibrium given by equation (54), derived in the approximation of $\Delta\beta \ll (a/h)^2$. Thus self-consistency implies that the validity of (62) is limited to $h/a \ll 0.13$. This condition is satisfied, albeit marginally, for the parameters discussed in §2. The instability threshold for more general equilibria, with higher values of h/a , is at present unknown.

8. MASS OF CONFINED ACCRETED MATTER

Assuming that the crust is perfectly rigid, the amount of mass needed to drive the instability is, from inequality (62),

$$\begin{aligned} \Delta M &= \pi a^2 \frac{B^2}{8\pi} \frac{\Delta\beta}{g} \\ &= 0.73 B^2 a^3 \frac{\rho}{p}. \end{aligned} \quad (63)$$

Substituting Eq. (2) in Eq. (63), we have

$$\begin{aligned} \Delta M &= 3.8 \times 10^{-13} M_\odot \left(\frac{B}{10^{12} \text{ G}} \right)^2 \left(\frac{a}{10^5 \text{ cm}} \right)^3 \times \\ &\quad \left(\frac{Z}{26} \right)^{2/3} \left(\frac{A}{56} \right) \left(\frac{5 \times 10^8 \text{ K}}{T} \right) \left(\frac{173}{\Gamma} \right). \end{aligned} \quad (64)$$

At a given accretion rate $\dot{M}$, the time necessary to accumulate this mass is $\Delta M/\dot{M} = 33 \text{ hr} (10^{-10} M_\odot \text{ yr}^{-1}/\dot{M})$. At present we cannot say what, if any, are the observable effects of the instability, but the timescale to accumulate a critical mass ΔM is well-suited for monitoring.

9. DISCUSSION

Using the MHD energy principle, we have shown that the accreted material on the polar cap of a strongly magnetized neutron star is subject to the ballooning instability when the overpressure in the vicinity of the crust exceeds the magnetic pressure by the factor $7.8a/h$, where a is the horizontal pressure gradient length scale (identified with the polar cap radius) and h is the density scale height. For typical polar cap parameters, the critical overpressure is $\sim 10^{-4}$ of the total plasma pressure. The instability is localized to within a density scale height of the field line footpoint (at the top of the crust) which implies that plasma is perturbed at densities $(\rho/\rho_0) > 0.33$; for ρ_0 found from equation (1) for a model atmosphere (cf. Brown & Bildsten 1999), with a temperature of $7.6 \times 10^8 \text{ K}$, assuming a locally Eddington accretion rate. This implies $\rho > 10^{10} \text{ g cm}^{-3}$. This density is higher than where the accreted hydrogen and helium ignite which suggests

that the ballooning instability near the threshold for onset will not affect the nuclear burning. It is not clear, however, whether the instability can degrade the confinement enough to prevent a further accumulation of mass well above the instability threshold.

In our stability analysis we have omitted the contribution of the Schwarzschild term to the MHD potential energy arguing that it is small in highly degenerate plasmas of uniform composition immersed in strong magnetic fields. To test the validity of this approximation we have performed a calculation along the same lines as in §4 and §5, including the buoyancy contribution, for $\epsilon = 4 \times 10^{-4}$, for a realistic atmosphere (cf. Brown & Bildsten 1999). We found that this inclusion increases the numerical factor on the RHS of inequality (62) by 17% for $B = 10^{12}$ G and by less than 1% for $B \geq 5 \times 10^{12}$. Thus the effect of buoyancy is indeed weak for magnetic fields of interest, as long as ϵ is as small as assumed.

The latter assumption, as well as that of composition uniformity, is violated in the electron-capture layers. The location of these layers is not precisely known. Haensel & Zdunik (1990) found that electron capture layers closest to the top of the crust occur at densities 10^{10} g cm $^{-3}$ and 8×10^{10} g cm $^{-3}$. Since the instability found in the present paper is localized within a density scale height from the crust which for the model atmosphere implies localization between densities 10^{10} and 3×10^{10} g cm $^{-3}$, we expect that the electron capture effects would not significantly modify our conclusions.

In our analysis the crust was modelled as a rigid boundary in which the magnetic field is anchored. For this approximation to be justified the lateral distortion of the crust in response to the overpressure must be much smaller than the magnetic field line distortion in the overlying fluid. This is true when $\Delta\beta/\beta$ exceeds $\alpha_s\Delta a/a$ where

$\Delta a \sim (h^2/a)\Delta\beta$ is the displacement of flux surfaces in the ocean (cf. §6), i.e., when $\beta > (a/h)^2/\alpha_s$; here $\alpha_s p$ is the crust shear modulus. Estimates (Pandharipande et al. 1976; Strohmayer et al. 1991) find $\alpha_s \sim 10^{-3} - 10^{-2}$, which for the characteristic density scale height and polar cap radius (cf. §1) implies that $\beta \gtrsim 10^4 - 10^5$. For the model atmosphere, this condition is satisfied for $B \lesssim 5 \times 10^{12}$ G.

The imperfect rigidity of the crust may affect the estimate of the accreted mass necessary to destabilize the ballooning mode (eq. [63]). Ushomirsky et al. (2000) found that the crust tended to sink in response to a lateral density discontinuity. While their findings are not directly applicable to our case, it is nevertheless conceivable that the crust could deform sufficiently to significantly reduce the applied overpressure. The mass required to drive the instability would then be higher than our estimate.

The effect of the instability is an enhanced cross-field transport of matter and a resulting decrease in the plasma confinement time in the polar cap. To find the magnitude of this anomalous transport requires analysis of nonlinear effects which are beyond the scope of this paper. The analysis of these effects and, in particular, the question of the existence of a steady state, in which the transport is sufficiently rapid to balance the influx of the accreted matter, are relegated to future publications.

The authors thank Ellen Zweibel for pointing out the potential importance of the stabilizing effect of the Schwarzschild term in equation (21). Helpful discussions with Thierry Emonet, Timur Linde, and Dmitri Uzdensky are also acknowledged. This research was supported by the ASCI Center for Astrophysical Thermonuclear Flashes at the University of Chicago under Department of Energy contract B341495. EFB acknowledges support from an Enrico Fermi Fellowship at the University of Chicago.

REFERENCES

- Arons, J. & Lea, S. M. 1980, *ApJ*, 235, 1016
 Basko, M. M. & Sunyaev, R. A. 1976, *MNRAS*, 175, 395
 Bernstein, I.B., Frieman, I.A., Kruskal, M.D. & Kulsrud, R.M., 1958, *Proc. R. Soc. (London) A*, 224, 17
 Bildsten, L. 1998, in *The Many Faces of Neutron Stars*, ed. A. Alpar, R. Buccheri, & J. van Paradijs, Vol. 515, NATO ASI ser. C (Dordrecht: Kluwer), 419
 Bildsten, L., Chakrabarty, D., Chiu, J., Finger, M. H., Koh, D. T., Nelson, R. W., Prince, T. A., Rubin, B. C., Scott, D. M., Stollberg, M., Vaughan, B. A., Wilson, C. A., & Wilson, R. B. 1997, *ApJS*, 113, 367
 Brown, E. F. & Bildsten, L. 1998, *ApJ*, 496, 915
 Connor, J. W., Hastie, R. J. and Taylor, J. B., 1979, *Proc. R. Soc. (London) A* 365, 1
 Coppi, B., & Rosenbluth, M. N., 1966, *Plasma Physics and Controlled Fusion Research 1965* (IAEA: Vienna)
 Davidson, K. & Ostriker, J. P. 1973, *ApJ*, 179, 585
 De Bruyne, P., Hood, A. W., 1989, *Solar Phys.*, 123, 241
 Elsner, R. F. & Lamb, F. K. 1977, *ApJ*, 215, 897
 Farouki, R. & Hamaguchi, S. 1993, *Phys. Rev. E*, 47, 4330
 Freidberg, J.P., 1987, *Ideal Magnetohydrodynamics* (Plenum Press: New York)
 Furth, H. P., Killeen, J., Rosenbluth, M. N., & Coppi, B., 1965, in *Plasma Physics and Controlled Thermonuclear Research 1964* (IAEA: Vienna)
 Greene, J. M., & Johnson, J. L., 1968, *Plasma Phys.*, 10, 729
 Haensel, & P., Zdunik, J. L., 1990, *Astron. Astrophys.*, 229, 117
 Hameiri, E., Laurence, P., Mond, M., 1991, *J. Geophys. Res.*, 93, 1513
 Hameury, J. M., Bonazzola, S., Heyvaerts, J., & Lasota, J. P. 1983, *A&A*, 128, 369
 Heindl, W. A., Coburn, W., Gruber, D. E., Pelling, M., Rothschild, R. E., Kretschmar, P., Kreykenbohm, I., Wilms, J., Pottschmidt, K., & Staubert, R. 2000, in "Proceedings of the 5th Compton Symposium", in press
 Hood, A. W., 1986, *Solar Phys.*, 103, 329
 Inogamov, N. A. & Sunyaev, R. A. 1999, *Astronomy Letters*, 25, 269
 Joss, P. C. & Li, F. K. 1980, *ApJ*, 238, 287
 Kruskal, M.D. & Schwarzschild, M., 1954, *Proc. R. Soc. (London) A*, 223, 63
 Lamb, F. K., Pethick, C. J., & Pines, D. 1973, *ApJ*, 184, 271
 Lewin, W. H. G., van Paradijs, J., & Taam, R. 1995, in *X-ray Binaries*, ed. W. H. G. Lewin, J. van Paradijs, & E. P. J. van den Heuvel (Cambridge University Press), 175–232
 McNutt, R. L., Coppi, P. S., Selesnick, R. S., & Coppi, B., 1987, *J. Geophys. Res.*, 92, 4377
 Ohkawa, T., Kerst, D. W., 1961, *Nuovo Cim.*, 22, 784
 Pandharipande, V. R., Pines, D., & Smith, R. A. 1976, *ApJ*, 208, 550
 Parker, E. N., 1966, *ApJ* 145, 811
 Popham, R. & Sunyaev, R. 2000, *ApJ*, submitted. Preprint available: astro-ph/0004017
 Pringle, J. E. & Rees, M. J. 1972, *A&A*, 21, 1
 Schatz, H., Bildsten, L., Cumming, A., & Wiescher, M. 1999, *ApJ*, 524, 1014
 Shibata, K., Tajima, T., Matsumoto, R., Horiuchi, T., Hanawa, T., Rosner, R., Uchida, Y., *Astrophys. J.*, 338, 471
 Strauss, H. R., Longcope, D. W., *Solar Phys.* 149, 63
 Strohmayer, T., van Horn, H. M., Ogata, S., Iyetomi, H., & Ichimaru, S. 1991, *ApJ*, 375, 679
 Ushomirsky, G., Cutler, C., & Bildsten, L. 2000, *MNRAS*, submitted
 White, N. E., Nagase, F., & Parmar, A. N. 1995, in *X-ray binaries*, ed. W. H. G. Lewin, J. van Paradijs, & E. P. J. van den Heuvel (Cambridge: Cambridge University Press), 1–57